

Kaposiform intramuscular hemangiosarcoma in a hoary fox (*Lycalopex vetulus*)¹

Davi S. Leal^{2*}, Daniela O. Rocha², Gabriel A.O. Bezerra², Isabel Luana de Macêdo², Liz A. Cerqueira²,
Dara E. V. da Costa², Líría Q. L. Hirano² and Márcio B. Castro²

Registros ORCID:

Davi de Souza Leal – <https://orcid.org/0000-0001-9547-3633>

Daniela Oliveira Rocha – <https://orcid.org/0000-0002-6515-6992>

Gabriel Almeida de Oliveira Bezerra – <https://orcid.org/0009-0000-8653-7332>

Isabel Luana de Macêdo – <https://orcid.org/0000-0001-7327-9143>

Liz de Albuquerque Cerqueira – <https://orcid.org/0000-0002-6353-6238>

Dara E. V. da Costa – <https://orcid.org/0000-0002-9329-0339>

Líría Q. L. Hirano – <https://orcid.org/0000-0002-9438-7538>

Márcio Botelho de Castro – <https://orcid.org/0000-0002-3996-0856>

ABSTRACT.- Leal D.S., Rocha D.O., Bezerra G.A.O., Macêdo I.L., Cerqueira L.A., Costa D.E. V., Hirano L.Q. L. & Castro M.B. **Kaposiform intramuscular hemangiosarcoma in a hoary fox (*Lycalopex vetulus*)**. *Pesquisa Veterinária Brasileira* 46:e07859, 2026. Laboratório de Patologia e Perícia Veterinária, Faculdade de Agronomia e Medicina Veterinária (FAV), Campus Universitário Darcy Ribeiro, Universidade de Brasília (UnB), Brasília, DF, Brasil. E-mail: mbcastro@unb.br

Hemangiosarcoma (HSA) is a malignant vascular neoplasm commonly reported in domestic dogs but rarely described in wild canids. The hoary fox (*Lycalopex vetulus*), a near-threatened Neotropical canid endemic to the Brazilian Cerrado, has rarely been associated with neoplastic disease. This report describes the clinical, gross, histopathological, and immunohistochemical findings of a Kaposiform intramuscular HSA in a captive adult male hoary fox from Brazil. Clinically, the animal presented with moderate lameness and swelling of the left pelvic limb. Grossly, a poorly demarcated, infiltrative mass involved the proximal femoral musculature and showed irregular white, red, and black areas. Histologically, the neoplasm consisted of spindle-shaped cells arranged predominantly in solid sheets, with occasional irregular vascular channels, and was associated with marked pleomorphism and nineteen mitotic figures in 10 high-power microscopic fields (2.37 mm²). Immunohistochemically, neoplastic cells were positive for CD31, Factor VIII, and vimentin. To our knowledge, this is the first report of Kaposiform intramuscular HSA in a Neotropical wild canid.

INDEX TERMS: neoplasm, wild canids, immunohistochemistry, Neotropical canids, oncology.

RESUMO.- [Hemangiossarcoma Kaposiforme intramuscular em uma raposa-do-campo (*Lycalopex vetulus*).] O hemangiossarcoma (HSA) é uma neoplasia vascular maligna frequentemente descrita em cães domésticos, mas raramente em canídeos silvestres. A raposinha do campo (*Lycalopex vetulus*), um canídeo neotropical praticamente ameaçado e endêmico do Cerrado brasileiro, raramente tem sido associada a doenças neoplásicas. Este relato descreve os achados clínicos, macroscópicos, histopatológicos e imuno-histoquímicos de um HSA intramuscular do subtipo Kaposiforme em um macho adulto cativo de *Lycalopex vetulus* no Brasil. Clinicamente, o animal apresentava claudicação e aumento de volume moderados no membro pélvico esquerdo. Macroscopicamente, observou-se uma massa infiltrativa pouco delimitada envolvendo a musculatura femoral proximal, e apresentava áreas irregulares brancas, vermelhas e pretas. Histologicamente, a neoplasia era composta por células fusiformes, dispostas predominantemente em mantos sólidos, com canais vasculares irregulares ocasionais, associadas a acentuado pleomorfismo e dezenove figuras mitóticas em 10 campos de grande aumento microscópico (2,37 mm²). No exame imuno-histoquímico, as células neoplásicas apresentaram

¹ Received on may 28, 2026.

Accepted for publication on October 26, 2026.

² Laboratório de Patologia e Perícia Veterinária, Faculdade de Agronomia e Medicina Veterinária (FAV), Campus Universitário Darcy Ribeiro, Universidade de Brasília (UnB), Brasília, DF, Brasil. E-mail: mbcastro@unb.br

imunomarcção positiva para CD31, fator VIII e vimentina. Até o momento, este é o primeiro relato de HSA intramuscular Kaposiforme em um canídeo silvestre Neotropical.

TERMOS DE INDEXAÇÃO: neoplasia, canídeos selvagens, imunohistoquímica, canídeos neotropicais, oncologia.

INTRODUCTION

Neoplasms have been reported sporadically in wild canids, typically affecting aged, captive individuals (Lamglait et al., 2015; Acton et al., 2000). In contrast, in a study of more than 100,000 domestic dog tissue samples submitted for histopathology in Germany, neoplasms accounted for 64.7% of diagnoses (Aupperle-Lellbach et al., 2022). The hoary fox (*Lycalopex vetulus*) is a near-threatened Neotropical wild canid (Lemos et al., 2020) from the Brazilian Cerrado (Brazilian Savannah). Neoplasms are rarely reported in this species, with one described case of a metastatic oral carcinoma in a female (Zanoti et al., 2024).

Hemangiosarcomas (HSA) are malignant neoplasms arising from vascular endothelial cells, characterized by highly infiltrative behavior and marked metastatic potential (Roccabianca et al., 2020; Suzuki et al., 2024). In domestic dogs, these neoplasms most frequently involve the spleen, liver, heart, and skin (De Nardi et al., 2023), although musculoskeletal presentations have also been reported (Carloni et al., 2019). However, the occurrence, anatomical distribution, and biological behavior of HSA remain poorly documented in wild canids (Newell-Fugatea et al., 2009; Lamglait et al., 2015; Sarvi et al., 2024; Sato et al., 2024).

Epidemiological data from domestic animals show a substantially higher frequency of neoplasms diagnosed by necropsy and biopsy than that reported in wildlife studies. In a series of 12,118 domestic animal specimens submitted for pathological evaluation, 16% were diagnosed with neoplastic lesions, predominantly in dogs (Kimura et al. 2012). This contrasts with the 2.75% prevalence of neoplasms reported in captive wild mammals at the San Diego Zoo (Effron et al. 1977).

Neoplasms are a significant cause of morbidity and mortality in domestic dogs, particularly in geriatric animals, with HSA accounting for approximately 5% of primary non-cutaneous malignant neoplasms in dogs (Griffin et al., 2021). In contrast, information on neoplastic diseases in wild canids remains scarce and is derived primarily from retrospective studies of captive animals in zoological collections. Although several neoplasms have been described in African wild dogs (*Lycaon pictus*), including carcinomas, soft tissue sarcomas, mammary and uterine tumors, and cutaneous apocrine neoplasms, reports of HSA remain rare (Lamglait et al., 2015; Mitchell et al., 2023). Similarly, in Neotropical canids, such as the maned wolf (*Chrysocyon brachyurus*), the most frequently reported neoplasms include ovarian dysgerminomas and mammary carcinomas, as well as a rare case of metastatic oral squamous cell carcinoma in a hoary fox (Marcinczyk et al., 2024; Zanoti et al., 2024). Furthermore, unspecified musculoskeletal sarcomas have been reported in a survey of 171 captive maned wolves in zoological collections (Marcinczyk et al., 2024).

Given the limited understanding of neoplasms in Neotropical wild canids, documenting and characterizing underreported neoplasms in wildlife species can elucidate causes of morbidity and contribute to conservation medicine for these species. Herein, we describe a case of Kaposiform intramuscular hemangiosarcoma in a captive hoary fox (*Lycalopex vetulus*), highlighting the clinical, gross, histopathological, and immunohistochemical findings.

MATERIALS AND METHODS

Ethical approval. In this study, the only material analyzed was tissue samples submitted for routine pathological evaluation. Therefore, no animal experiments were conducted, and ethical approval was not required under the guidelines of the National Council for the Control of Animal Experimentation (CONCEA). The data supporting the findings of this study are available from the corresponding author upon reasonable request.

According to the clinical history report, an adult male hoary fox (*Lycalopex vetulus*) weighing 4.5 kg from the Brasília Zoological Garden Foundation, Brazil, presented with moderate lameness of the left pelvic limb and swelling in the proximal femoral region. A fine-needle aspiration biopsy was performed, and the cytological diagnosis was suggestive of a malignant spindle cell neoplasm. The animal underwent amputation of the left pelvic limb via coxofemoral disarticulation, and the specimen was sent for pathological evaluation to the Veterinary and Forensic Pathology Laboratory of the University of Brasília, Brazil.

The amputated limb was received with intact, haired skin and showed marked enlargement of the proximal femoral region (Fig. 1). Tissue samples were collected, fixed in 10% neutral-buffered formalin, routinely processed, and stained with hematoxylin and eosin (H&E) for histological examination.

Additionally, tumor tissue samples were submitted for immunohistochemical analysis using the biotin-streptavidin-peroxidase method with 3,3'-diaminobenzidine (DAB), 4 min (ImmunoDetector DAB, HRP, BioSB Inc.). Antibodies against Factor VIII (dilution 1:400, Factor VIII; clone F8/86, Dako Corp.), CD31 (dilution 1:100, CD31, clone JC/70A, Dako Corp.), and vimentin (dilution 1:400, VIM, clone Vim 3B4, Dako Corp.) were applied to histological sections after antigen retrieval in a pressure cooker, followed by an overnight incubation. Positive

controls included a case of cutaneous canine hemangiosarcoma. Negative controls were incubated with normal rabbit serum (1:10 dilution) instead of the primary antibodies.

RESULTS

Gross findings included a heterogeneous, firm mass measuring 5.9 × 4.4 × 3.4 cm, with irregular white, red, and black areas. The mass expanded and infiltrated the tensor fasciae latae, vastus lateralis, semitendinosus, and semimembranosus muscles (Fig. 1).

Microscopically, skeletal muscle bundles were infiltrated by a spindle cell proliferation arranged in solid bundles and occasionally forming irregular vascular spaces (Fig. 2A). The neoplastic cells had indistinct cell borders and eosinophilic cytoplasm (Fig. 2B). The nuclei were oval, with coarse chromatin and one to two prominent nucleoli. Nineteen mitotic figures were observed in 10 high-power microscopic fields (2.37 mm²). Extensive areas of necrosis, hemorrhage, and edema were interspersed within the neoplastic proliferation.

Neoplastic cells showed moderate immunolabeling for Factor VIII (Fig. 3A), marked immunostaining for CD31 (Fig. 3B), and VIM. Based on histopathological and immunohistochemical findings, the neoplasm was classified as a Kaposiform intramuscular HSA, exhibiting morphological features consistent with those described for this subtype in canine HSAs (Roccabianca et al., 2020; Suzuki et al., 2024).

DISCUSSION

Vascular neoplasms, particularly hemangiosarcomas (HSAs), are rarely reported in wild canids from the Northern Hemisphere and Africa, with sporadic cases involving visceral organs such as the right atrium, spleen, and liver (Newell-Fugate et al., 2009; Sarvi et al., 2024). To date, only one metastatic case involving the thoracic musculature has been described in a red wolf (*Canis rufus*) (Sarvi et al., 2024). In contrast, HSAs have not yet been reported in Neotropical wild canids to the authors' knowledge.

The lameness observed in this case is consistent with the clinical presentation of musculoskeletal HSA in domestic dogs. In a previous study, 9 of 15 (60%) dogs with muscular HSA metastases exhibited lameness or reluctance to move, often without other systemic clinical signs, leading to initial presumptive diagnoses of primary orthopedic or neurological disorders (Carloni et al., 2019). Such presentations may contribute to delayed diagnosis, particularly because deep intramuscular masses can clinically resemble inflammatory conditions and other non-neoplastic lesions, such as abscesses, hematomas and granulomatous inflammations (Gasparini et al., 2020).

In the present case, the infiltrative, poorly demarcated appearance of the intramuscular neoplasm hindered initial suspicion of a neoplastic process, as no externally visible nodular masses were observed, unlike the exophytic lesions typically associated with cutaneous HSAs (Carloni et al., 2019). On cut section, the lesion in the hoary fox exhibited a heterogeneous appearance with extensive hemorrhagic, white, and black areas, suggesting necrosis, consistent with descriptions of HSAs in both domestic (De Nardi et al., 2023) and wild canids (Kim et al., 2015; Sarvi et al., 2024). These features are likely related to tortuous, irregular vascular channels that are frequently obstructed by thrombi, leading to ischemia, hemorrhage, and tissue necrosis. HSAs may expand locally through tissue infiltration or metastasize to distant organs and can present as solitary, multifocal, or disseminated lesions (Suzuki et al., 2024). Because intramuscular HSAs generally lack the reddish or purplish appearance characteristic of cutaneous vascular tumors, they may be easily confused with other neoplastic or non-neoplastic conditions (De Nardi et al., 2023).

The Kaposiform intramuscular HSA in the hoary fox showed a predominantly microscopic solid growth pattern, with sparse irregular vascular channels and numerous mitotic figures, findings consistent with poorly differentiated forms of this neoplasm (Roccabianca et al., 2020; Suzuki et al., 2024). In domestic dogs, HSA is recognized as a histologically heterogeneous tumor, ranging from well-formed anastomosing vascular channels to solid sheets of pleomorphic endothelial cells (Roccabianca et al., 2020; De Nardi et al., 2023). Unfortunately, in wild canids such as African wild dogs and red and gray wolves (Newell-Fugate et al., 2009; Sarvi et al., 2024), histological classification of HSAs has not been performed for comparison with the neoplasm in the hoary fox.

In the present case, the absence of post-surgical follow-up limited further clinical evaluation. Although histological parameters such as differentiation, mitotic index, size, and invasiveness are prognostic indicators in canine soft tissue sarcomas (Kuntz et al., 1997), their relevance for hemangiosarcomas remains unvalidated, with no clear association with survival or clinical outcome (Schultheiss et al., 2004; Bertazzolo et al., 2005). Furthermore, the lack of an established histological grading system for HSA (Bertazzolo et al., 2005; Shiu et al., 2011), together with the scarcity of reports in wild canids (Newell-Fugate et al., 2009; Sarvi et al., 2024), precludes reliable prognostic assessment in this hoary fox.

To the best of our knowledge, this is the first report of a Kaposiform intramuscular HSA in a neotropical canid. However, it was not possible to determine whether the lesion was a primary intramuscular neoplasm or a metastatic focus, as only the amputated limb was available for evaluation, precluding assessment of distant metastatic sites and identification of a potential primary tumor. In dogs, muscular involvement by HSA is frequently associated with advanced disseminated disease affecting multiple organs (Carloni et al., 2019), although this could not be assessed in the hoary fox. A comparable case has been described in a male red wolf with

metastatic HSA infiltrating the intercostal musculature of the right lateral thorax from the adjacent subcutaneous tissue (Sarvi et al., 2024). Nevertheless, visceral forms remain the most commonly reported presentations of HSA in wild canids, particularly involving the spleen, liver, and heart (Newell-Fugate et al., 2009; Lamglait et al., 2015; Sarvi et al., 2024; Sato et al., 2024; Minichino et al., 2026).

Given the histopathological features of the Kaposiform intramuscular HSA observed in this study, immunohistochemistry was essential for definitive diagnosis, with neoplastic cells showing immunolabeling for vimentin and the endothelial markers CD31 and Factor VIII-related antigen. In spindle-cell neoplasms with predominantly solid growth patterns, specific endothelial markers are critical to confirm vascular origin (Bertazzolo et al., 2005; Chu et al., 2023) and to differentiate HSA from other primary or infiltrative intramuscular neoplasms, such as fibrosarcoma, schwannoma, and hemangiopericytoma. These neoplasms may display spindle-cell or anaplastic morphological features similar to those of solid HSAs but are typically negative for CD31 and Factor VIII immunolabeling (Ferrer et al., 1995).

The principal differential diagnoses for intramuscular hemangiosarcoma include hematomas, abscesses, hemangiomas, other spindle-cell soft-tissue sarcomas, and, in epithelioid variants, metastatic undifferentiated carcinomas (Russell et al., 1977; Sabattini & Bettini, 2009). Among soft-tissue sarcomas, fibrosarcoma, leiomyosarcoma, and rhabdomyosarcoma may exhibit spindle-cell morphology and infiltrative growth within skeletal muscle; however, they differ from hemangiosarcoma by the absence of vascular channel formation and by distinct immunohistochemical profiles (Russell et al., 1977; Shi et al., 2023). In addition, hematomas and other soft-tissue neoplasms (Olimpo et al., 2023; Caraffi et al., 2025), injection-site sarcomas (Jacobs et al., 2017), and foreign-body-associated sarcomas, including those related to microchips (Vascellari et al., 2006), may further complicate the definitive diagnosis of HSA. These were excluded herein based on the clinical history and pathological findings in the present case.

Positive immunolabeling for endothelial markers, together with histopathological features, was essential for confirming the vascular endothelial origin of the neoplasm and for excluding other nonvascular sarcomas in the hoary fox. Despite the study's limitations, this report contributes to current knowledge of diseases associated with lameness in Neotropical canids and expands the pathological characterization of a rare neoplasm affecting skeletal muscle.

The most relevant limitations of this study included restricted clinical information, no data regarding hematological or biochemical findings, therapeutic management, postoperative follow-up, or survival could be retrieved despite efforts to obtain further information. In addition, because only the amputated limb was available for pathological evaluation, complete tumor staging and assessment of distant metastases were not possible, precluding determination of whether the lesion represented a primary intramuscular neoplasm or metastatic disease.

CONCLUSION

This report documents the first known case of Kaposiform intramuscular HSA in a hoary fox, highlighting the diagnostic challenges posed by intramuscular HSAs, particularly when clinical signs are nonspecific (lameness) and external masses are absent. Histopathology combined with immunohistochemistry was essential for establishing the definitive diagnosis and differentiating the tumor from other spindle-cell neoplasms of skeletal muscle. Although the absence of complete staging and clinical follow-up limited assessment of the tumor's biological behavior and prognosis, the findings suggest an aggressive and infiltrative neoplasm. Given the scarcity of information on neoplastic diseases in Neotropical canids, this report reinforces the importance of including HSA in the differential diagnosis of muscular lesions associated with locomotor abnormalities in captive wildlife species.

Conflict of interest statement.- The authors declare that they have no conflicts of interest.

Acknowledgments.- We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Brazil, for partially funding the doctoral scholarship of Isabel L. Macêdo Liz and C. Albuquerque (Finance Code 001). Special thanks are extended to the National Council for Scientific and Technological Development (CNPq) for financial support (Grant No. 307909/2021-2 to M.B. Castro).

CRedit author statement.- Davi S. Leal and Márcio B. Castro: Conceptualization, Supervision, Project administration, Resources, , Writing – original draft, Writing – review & editing. Liz A. Cerqueira: Data curation, Investigation, Visualization, Writing – original draft, Writing – review & editing. Daniela O. Rocha, Gabriel A.O. Bezerra, Isabel Luana de Macêdo, Dara E. V. da Costa, and Líria Q. L. Hirano: Data curation, Formal analysis, Investigation, Methodology, Writing – review & editing. All authors approved the final version of the manuscript.

Editor-in-Chief.- Fabiano José Ferreira de Sant'Ana.

REFERENCES

- Acton AE, Munson L, Waddell WT. Survey of necropsy results in captive red wolves (*Canis rufus*), 1992-1996. *J Zoo Wildl Med* 2000;31(1):2-8. [https://doi.org/10.1638/1042-7260\(2000\)031\[0002:SONRIC\]2.0.CO;2](https://doi.org/10.1638/1042-7260(2000)031[0002:SONRIC]2.0.CO;2)
- Aupperle-Lellbach H, Grassinger JM, Floren A, Törner K, Beitzinger C, Loesenbeck G, et al. Tumour incidence in dogs in Germany: a retrospective analysis of 109,616 histopathological diagnoses (2014-2019). *J Comp Pathol* 2022;198:33-55. <https://doi.org/10.1016/j.jcpa.2022.07.009>
- Bertazzolo W, Dell'Orco M, Bonfanti U, Ghisleni G, Caniatti M, Masserdotti C, et al. Canine angiosarcoma: cytologic, histologic, and immunohistochemical correlations. *Vet Clin Pathol* 2005;34(1):28-34. <https://doi.org/10.1111/j.1939-165X.2005.tb00005.x>
- Carafi OA, O'Sullivan J, Hoey S, Kelly PA, Mullins RA. Intramuscular hemophilic pseudotumor in a dog: imaging findings and surgical treatment. *Vet Radiol Ultrasound* 2025;66(5):e70071. <https://doi.org/10.1111/vru.70071>
- Carloni A, Terragni R, Morselli-Labate AM, Paninarova M, Graham J, Valenti P, et al. Prevalence, distribution, and clinical characteristics of hemangiosarcoma-associated skeletal muscle metastases in 61 dogs: a whole body computed tomographic study. *J Vet Intern Med* 2019;33(2):812-819. <https://doi.org/10.1111/jvim.15456>
- Chu KT, Nekouei O, Sandy JR. Histopathological grading, clinical staging and CD31 expression of canine splenic hemangiosarcoma. *Vet Sci* 2023;10(3):190. <https://doi.org/10.3390/vetsci10030190>
- De Nardi AB, De Oliveira Massoco Salles Gomes C, Fonseca-Alves CE, De Paiva FN, Linhares LCM, Carra GJU, et al. Diagnosis, prognosis, and treatment of canine hemangiosarcoma: a review based on a consensus organized by the Brazilian Association of Veterinary Oncology, ABROVET. *Cancers (Basel)* 2023;15(7):2025. <https://doi.org/10.3390/cancers15072025>
- Effron, M., Griner, L., & Benirschke, K. (1977). Nature and rate of neoplasia found in captive wild mammals, birds, and reptiles at necropsy. *Journal of the National Cancer Institute*, 59(1), 185-198. <https://doi.org/10.1093/jnci/59.1.185>
- Ferrer L, Fondevila D, Rabanal RM, Vilafranca M. Immunohistochemical detection of CD31 antigen in normal and neoplastic canine endothelial cells. *J Comp Pathol* 1995;112(4):319-326. [https://doi.org/10.1016/S0021-9975\(05\)80013-1](https://doi.org/10.1016/S0021-9975(05)80013-1)
- Gasparini MK, Selmic LE, Cook M, Brown M, Wavreille VA. Abscessation associated with an intramuscular haemangiosarcoma in a dog. *Vet Rec Case Rep* 2020;8:e000982. <https://doi.org/10.1136/vetreccr-2019-000982>
- Griffin MA, Culp WTN, Rebhun RB. Canine and feline haemangiosarcoma. *Vet Rec* 2021;189(9):e585. <https://doi.org/10.1002/vetr.585>
- Jacobs TM, Poehlmann CE, Kiupel M. Injection-site sarcoma in a dog: clinical and pathological findings. *Case Rep Vet Med* 2017;2017:6952634. <https://doi.org/10.1155/2017/6952634>
- Kim JH, Graef AJ, Dickerson EB, Modiano JF. Pathobiology of hemangiosarcoma in dogs: research advances and future perspectives. *Vet Sci* 2015;2(4):388-405. <https://doi.org/10.3390/vetsci2040388>
- Kimura KC, Gárate AP, Dagli MLZ. Retrospective study of neoplasms in domestic animals: a survey between 1993 and 2002 of the Service of Animal Pathology, Department of Pathology, School of Veterinary Medicine and Animal Science, University of Sao Paulo, Southeast Brazil. *Braz J Vet Pathol* 2012;5(2):60-69. <https://doi.org/10.24070/bjvp.1983-0246.005011>
- Kuntz CA, Dernell WS, Powers BE, Devitt C, Straw RC, Withrow SJ. Prognostic factors for surgical treatment of soft-tissue sarcomas in dogs: 75 cases (1986-1996). *J Am Vet Med Assoc* 1997;211(9):1147-1151.
- Lamglait B, Trunet E, Leclerc A. Retrospective study of mortality of captive African wild dogs (*Lycaon pictus*) in a French zoo (1974-2013). *J Zoo Aquar Res* 2015;3(2):47-51. <https://doi.org/10.19227/jzar.v3i2.124>
- Lemos FG, Azevedo FC, Paula RC, Dalponte JC. *Lycalopex vetulus*. The IUCN Red List of Threatened Species 2020:e.T6926A87695615. <https://doi.org/10.2305/IUCN.UK.2020-2.RLTS.T6926A87695615.en>
- Marcinczyk M, Songsasen N, Hammond EE, DeCesare G, Rodden M, McAloose D. A retrospective study of disease processes in maned wolves (*Chrysocyon brachyurus*) in North American zoological institutions with emphasis on urolithiasis, inflammatory bowel disease, and neoplasia. *J Zoo Wildl Med* 2024;54(4):681-691. <https://doi.org/10.1638/2023-0024>
- Minichino A, Mennonna G, Dimatteo M, Degli Uberti B, D'Amore M, Santomartino F, et al. Case report: renal hemangiosarcoma in a free-ranging red fox (*Vulpes vulpes*). *Front Vet Sci* 2026;13:1766425. <https://doi.org/10.3389/fvets.2026.1766425>
- Mitchell EP, Henker MS, Lemberger K, Gohl C, Majzoub-Altweck M, Falkenau A, et al. Cutaneous apocrine gland neoplasia in 16 captive African wild dogs (*Lycaon pictus*). *J Comp Pathol* 2023;207:59-65. <https://doi.org/10.1016/j.jcpa.2023.10.005>
- Newell-Fugate A, Lane E. Periaortic haemangiosarcoma in an African wild dog (*Lycaon pictus*). *J S Afr Vet Assoc* 2009;80(2):108-110. <https://doi.org/10.4102/jsava.v80i2.182>
- Olimpo M, Buracco P, Ferraris EI, Piras LA, Maniscalco L, Giacobino D, et al. Surgical excision of intramuscular sarcomas: description of three cases in dogs. *Animals (Basel)* 2023;13(2):218.

<https://doi.org/10.3390/ani13020218>.

- Roccabianca, P., Schulman, F., Avallone, G., Dittmer, K., Kiupel, M., 2020. Malignant vascular tumors. In: Kiupel, M. (Ed.), *Surgical Pathology of Tumors of Domestic Animals: Tumors of Soft Tissue*, Vol. 3. Davis-Thompson Foundation, Gurnee, pp. 169–182.
- Russell WO, Cohen J, Enzinger F, Hajdu SI, Heise H, Martin RG, et al. A clinical and pathological staging system for soft tissue sarcomas. *Cancer* 1977;40(4):1562-1570. [https://doi.org/10.1002/1097-0142\(197710\)40:4<1562::aid-cnrcr2820400428>3.0.co;2-6](https://doi.org/10.1002/1097-0142(197710)40:4<1562::aid-cnrcr2820400428>3.0.co;2-6).
- Sabattini S, Bettini G. An immunohistochemical analysis of canine haemangioma and haemangiosarcoma. *J Comp Pathol* 2009;140:158-168. <https://doi.org/10.1016/j.jcpa.2008.10.006>
- Sarvi JY, Dowling KE, Swenson J, Haefele HJ, Garner MM. Hemangiosarcoma in red wolves (*Canis rufus*) and gray wolves (*Canis lupus*) in human care: six cases. *J Zoo Wildl Med* 2024;55(3):819-826. <https://doi.org/10.1638/2024-0007>
- Sato T, Nakamura SI, Kadekaru S, Endo M, Hattori YK, Une Y. Pathological findings of mammary gland carcinoma and hemangiosarcoma cases in two African wild dogs (*Lycaon pictus*). *J Vet Med Sci* 2024;86(5):596-599. <https://doi.org/10.1292/jvms.23-0381>.
- Schultheiss PC. A retrospective study of visceral and nonvisceral hemangiosarcoma and hemangiomas in domestic animals. *J Vet Diagn Invest* 2004;16(6):522-526. <https://doi.org/10.1177/104063870401600606>
- Shi J, Gao R, Zhang J, Xu R, Jia Q, Ma Y, et al. Invasive spindle-cell rhabdomyosarcoma with osteolysis in a dog: case report and literature review. *J Vet Diagn Invest* 2023;35(2):168-172. <https://doi.org/10.1177/10406387221147319>
- Shiu KB, Flory AB, Anderson CL, Wypij J, Saba C, Wilson H, Kurzman I, Chun R. Predictors of outcome in dogs with subcutaneous or intramuscular hemangiosarcoma. *J Am Vet Med Assoc* 2011;238(4):472-9. <https://doi.org/10.2460/javma.238.4.472>
- Suzuki, T., Henshaw, M. J., Yanagi, T., & Aoshima, K. Current understanding of comparative pathology and prospective research approaches for canine hemangiosarcoma. *Res Vet Sci*, 2024;167:105120. <https://doi.org/10.1016/j.rvsc.2023.105120>
- Vascellari M, Melchiotti E, Mutinelli F. Fibrosarcoma with typical features of postinjection sarcoma at site of microchip implant in a dog: histologic and immunohistochemical study. *Vet Pathol* 2006;43(4):545-548. <https://doi.org/10.1354/vp.43-4-545>.
- Zanoti IC, Gouveia BA, Ramos FR, Klein M, Momo C. Oral squamous cell carcinoma with lung and kidney metastases in a hoary fox (*Lycalopex vetulus*). *Braz J Vet Pathol* 2024;17(2):99-103. <https://doi.org/10.24070/bjvp.1983-0246.v17i2p99-103>

Figures



Fig.1. Kaposiform intramuscular hemangiosarcoma, hory fox (*Lycalopex vetulus*). Gross aspect. Amputated limb with an intact, marked swelling of the proximal femoral region. Inset: Cut surface. Heterogeneous mass with irregular white, red, and black areas infiltrating muscle tissues.




Fig.2. Kaposiform intramuscular hemangiosarcoma, hory fox (*Lycalopex vetulus*). Histological findings. **A.** Malignant solid proliferation composed of spindle cells arranged in bundles and occasionally forming vascular spaces (H&E, objective 10X). inset: bundles of malignant spindle cells (H&E, objective 40X). **B.** Spindle-shaped neoplastic cells, with moderate anisocytosis, anisocaryosis, and indistinct eosinophilic cytoplasm (H&E, objective 20X). inset: pleomorphic malignant cells forming vascular spaces (H&E, objective 40X).




Fig.3. Kaposiform intramuscular hemangiosarcoma, hory fox (*Lycalopex vetulus*). Immunohistochemical assay. **A.** Moderate immunolabeling of malignant endothelial cells for VWF (Immunoperoxidase, objective 20X). **B.** Spindle-shaped neoplastic cells with strong immunostaining for CD31 (Immunoperoxidase, objective 20X).

Online First